

ASSESSMENT OF PATHOGENIC OOMYCETES IMPACT ON *Salmosalar* L LARVAE USING OXIDATIVE STRESS BIOMARKERS

Eglė Gadeikytė^{1,2}, Gintarė Sauliūtė¹, Arvydas Markuckas², Milda Stankevičiūtė¹

¹Nature Research Centre, Akademijos St. 2, LT-08412 Vilnius, Lithuania

²Vilnius University, Life Sciences Center, Saulėtekio av. 7, 10223 Vilnius, Lithuania
egle.gadeikyte@gmc.stud.vu.lt

Freshwater fish are an important protein source for people in many countries, which is why aquaculture has now become a globally significant industry worldwide [1]. However, intensive aquaculture is related to the proliferation of parasites and other pathogenic organisms, posing a threat to biodiversity and food security worldwide [2]. One of these pathogens is oomycetes. Oomycetes, commonly known as water molds, are fungal-like microorganisms that can be parasitic towards a large number of plant and animal host species [3]. They cause one of the most destructive fish diseases in freshwater ecosystems – saprolegniasis [2]. Saprolegniasis is an infection that can develop at any stage of fish life [4] and is characterized mostly by a white or greyish cotton – wool like tuft found on infected skin, gills, or fish eggs. Due to the primary involvement of the skin in saprolegniasis, the disease is alternatively referred to as dermatomycosis.

Diseased fish in the most severe phase of infestation experience poor osmoregulation, respiratory failure, and, in certain cases, organ failure, which can lead to death [5]. It is critical to note that oomycete infections cause oxidative damage in fish, which contributes directly to disease pathogenesis [6]. Based on other research, *Saprolegnia parasitica* is the most important oomycete affecting freshwater fishes [6]. However, other oomycete species are also responsible for infestations, causing economically significant losses. For instance, *Saprolegnia australis*, acting as a pathogen on embryos and fry of salmonids, could colonize and cause their death [7].

The purpose of this research was to investigate the effects of *Saprolegnia* genus oomycetes on *Salmo salar* L. larvae. To achieve this aim, we evaluated the changes in enzyme glutathione S-transferases (GST) activity and levels of metallothionein (MTs). GST plays an important role in aquatic organisms protection from peroxidative damage [8]. MTs are metal-binding proteins with the ability to eliminate reactive oxygen species and maintain metal homeostasis in organisms [9]. Based on the results of present experiment, changes in GST activity were not detected. However, significant changes in MTs level were detected in oomycete-treated *S. salar* larvae compared with the control group.

Acknowledgments This research was funded by the Research Council of Lithuania, Project No. S-MIP-21-10, MULTIS.

-
- [1] Podeti. Koteswar Rao. Economically important freshwater fishes infected with fungi causes EUS. World J Adv Res Rev 2023; 17:605–609.
- [2] Pavić D, Grbin D, Hudina S, et al. Tracing the oomycete pathogen *Saprolegnia parasitica* in aquaculture and the environment. Sci Rep 2022; 12:16646.
- [3] Pavić D, Miljanović A, Grbin D, et al. Identification and molecular characterization of oomycete isolates from trout farms in Croatia, and their upstream and downstream water environments. Aquaculture 2021; 540:736652.
- [4] Barde RD. Clinical and pathological investigations in ulcer disease of *Cyprinus carpio* caused by *Aeromonas hydrophila*. ijhs 2022:3519–3526.
- [5] Lone SA, Manohar S. *Saprolegnia parasitica*, A Lethal Oomycete Pathogen: Demands to be Controlled. JIMB 2018; 6
- [6] Baldissera MD, Souza CF, Abbad LB, et al. Oxidative stress in liver of grass carp *Ctenopharyngodon idella* naturally infected with *Saprolegnia parasitica* and its influence on disease pathogenesis. Comp Clin Pathol 2020; 29:581–586.
- [7] Rezinciuc S, Sandoval-Sierra J-V, Diéguez-Uribeondo J. Molecular identification of a bronopol tolerant strain of *Saprolegnia australis* causing egg and fry mortality in farmed brown trout, *Salmo trutta*. Fungal Biology 2014; 118:591–600.
- [8] Park JC, Hagiwara A, Park HG, Lee J-S. The glutathione S-transferase genes in marine rotifers and copepods: Identification of GSTs and applications for ecotoxicological studies. Marine Pollution Bulletin 2020; 156:111080.
- [9] Kim J-H, Kang J-C. Oxidative stress, neurotoxicity, and metallothionein (MT) gene expression in juvenile rock fish *Sebastes schlegelii* under the different levels of dietary chromium (Cr6+) exposure. Ecotoxicology and Environmental Safety 2016; 125:78–84.