

CRISPR-CAS9 GENOME ENGINEERING IN *KLUYVEROMYCES MARXIANUS* FOR ENHANCED SECRETION OF RECOMBINANT ANTIBODIES

Justina Žičkutė^{1,2}, Danguolė Žiogienė², Alma Gedvilaitė²

¹Department of Biological DNA Modification, Institute of Biotechnology, Life Sciences Center, Vilnius University, Lithuania

²Department of Eukaryote Gene Engineering, Institute of Biotechnology, Life Sciences Center, Vilnius University, Lithuania
zickutejustina@gmail.com

Recombinant antibodies (RABs) are important in diagnostics, research, biotechnology, and therapeutics due to their high specificity, stability, and ease of modification [1]. Yeasts, which are easily genetically modified and cultivated, are often preferred as a cost-effective system for RABs production. The species *Kluyveromyces marxianus* is known for its efficient production and secretion of properly folded and active native and recombinant proteins, including RABs [2]. To improve RABs production technologies, yeasts can be genetically modified to create mutant strains with enhanced protein secretion properties.

Protein glycosylation, a crucial post-translational modification, involves attaching a glycan to a protein, ensuring its proper folding, activity, and stability. Dolichol kinase (DK), encoded by the essential *SEC59* gene, plays a role in glycosylation processes within the endoplasmic reticulum [3]. The reduced activity of DK, along with changes in glycosylation levels and the activity of other proteins in the secretory pathway, may lead to enhanced recombinant protein secretion [4]. Enhanced RABs secretion can also be achieved by reducing the activity of intracellular peptidases, as RABs are highly prone to proteolysis and often undergo degradation [5].

The aim of this study was to apply efficient CRISPR-Cas9 genome editing technology to construct a *K. marxianus* strain that displays improved secretion of RABs. In this investigation, a *K. marxianus* yeast WSS- $\Delta pep4$ strain was created by introducing mutations encoding G418S and I432S changes in the DK amino acids, and by disrupting the gene encoding vacuolar peptidase (*PEP4*). The introduction of mutations in the DK-encoding gene led to changes in DK activity, as indicated by reduced glycosylation efficiency of carboxypeptidase Y in the WSS strain. Additionally, the disruption of the *PEP4* gene in yeast resulted in a decrease in the proteolytic degradation of RABs. A secretion assay of the single-chain antibody fragment (scFv) linked to an antibody fragment crystallizable Fc (scFv-Fc) against *Gardnerella vaginalis* vaginolysin was performed and detected in yeast growth medium by Western blot. The results indicated that the constructed *K. marxianus* WSS- $\Delta pep4$ strain secreted recombinant scFv-Fc protein more efficiently compared to the wild-type *K. marxianus* strain. However, the secretion of RABs in yeast also depends on the specific properties of the recombinant protein, and further studies are necessary. The newly constructed *K. marxianus* WSS- $\Delta pep4$ mutant strain could be beneficial for future research aimed at enhancing RABs production technologies.

[1] Ma, H., O’Kennedy, R. (2017). Recombinant antibody fragment production. *Methods*, 116, 23–33.

[2] Nambu-Nishida, Y., Nishida, K., Hasunuma, T., Kondo, A. (2018). Genetic and physiological basis for antibody production by *Kluyveromyces marxianus*. *AMB Express*, 8(1), 56.

[3] Kale, D., Kikul, F., Phapale, P., Beedgen, L., Thiel, C., Brügger, B. (2023). Quantification of dolichyl phosphates using phosphate methylation and reverse-phase liquid chromatography–high resolution mass spectrometry. *Analytical Chemistry*, 95(6), 3210–3217.

[4] Žiogienė, D., Valavičiūtė, M., Norkienė, M., Timinskas, A., Gedvilaitė, A. (2019). Mutations of *Kluyveromyces lactis* dolichol kinase enhances secretion of recombinant proteins. *FEMS Yeast Research*, 19(3), foz024.

[5] Gast, V., Sandegren, A., Dunås, F., Ekblad, S., Güler, R., Thorén, S., Tous Mohedano, M., Molin, M., Engqvist, M. K. M., Siewers, V. (2022). Engineering *Saccharomyces cerevisiae* for the production and secretion of Affibody molecules. *Microbial Cell Factories*, 21(1), 36.